

Combinatorial Sums $\sum_{k \equiv r \pmod{m}} \binom{n}{k} a^k$ and Lucas Quotients (II)

Jiangshuai Yang

*Key Laboratory of Mathematics Mechanization, NCMIS, Academy of Mathematics
and Systems Science, Chinese Academy of Sciences, Beijing 100190, People's
Republic of China*
yangjiangshuai@amss.ac.cn

Yingpu Deng

*Key Laboratory of Mathematics Mechanization, NCMIS, Academy of Mathematics
and Systems Science, Chinese Academy of Sciences, Beijing 100190, People's
Republic of China*
dengyp@amss.ac.cn

Abstract

In [16], we obtained some congruences for Lucas quotients of two infinite families of Lucas sequences by studying the combinatorial sum

$$\sum_{k \equiv r \pmod{m}} \binom{n}{k} a^k.$$

In this paper, we show that the sum can be expressed in terms of some recurrent sequences with orders not exceeding $\varphi(m)$ and give some new congruences.

1. Introduction

Let p be an odd prime, using the formula for the sum

$$\sum_{k \equiv r \pmod{8}} \binom{n}{k},$$

Sun [12] proved that

$$\sum_{k=1}^{\frac{p-1}{2}} \frac{1}{k \cdot 2^k} \equiv \sum_{k=1}^{\lceil \frac{3p}{4} \rceil} \frac{(-1)^{k-1}}{k} \pmod{p}.$$

Later, Shan and E.T.H.Wang [5] gave a simple proof of the above congruence. In [10], Sun proved five similar congruences by using the formulas for Fibonacci quotient and Pell quotient.

In [13], Sun showed that the sum

$$\sum_{k \equiv r \pmod{m}} \binom{n}{k},$$

where n, m and r are integers with $m, n > 0$, can be expressed in terms of some recurrent sequences with orders not exceeding $\varphi(m)/2$, and obtained the following congruence

$$\sum_{k=1}^{\frac{p-1}{2}} \frac{3^k}{k} \equiv \sum_{k=1}^{\lfloor \frac{p}{6} \rfloor} \frac{(-1)^k}{k} \pmod{p}.$$

In [16], we studied more general sum

$$\sum_{k \equiv r \pmod{m}} \binom{n}{k} a^k, \tag{1}$$

and obtained congruences for Lucas quotients of two infinite families of Lucas sequences. See ([16] Theorems 4.10 and 5.4). In this paper, we continue studying the sum. We show that it can be expressed in terms of some recurrent sequences with orders not exceeding $\varphi(m)$, and obtain some new congruences.

For $x \in \mathbb{R}$, we use $[x]$ to denote the integral part of x i.e., the largest integer $\leq x$. For odd prime p and integer b , let $\left(\frac{b}{p}\right)$ denote the Legendre symbol and $q_p(b)$ denote the Fermat quotient $(b^{p-1} - 1)/p$ if $p \nmid b$. When $c, d \in \mathbb{Z}$, as usual (c, d) stands for the greatest common divisor of c and d . For any positive integer m , let $\zeta_m = e^{\frac{2\pi i}{m}}$ be the primitive m -th root of unity and let $\varphi(m)$, $\mu(m)$ denote the Euler totient function and Möbius function respectively. Throughout this paper, we fix $a \neq 0, \pm 1$.

2. Main Results

Definition 2.1. Let n, m, r be integers with $n > 0$ and $m > 0$. We define

$$\left[\begin{matrix} n \\ r \end{matrix} \right]_m(a) := \sum_{\substack{k=0 \\ k \equiv r \pmod{m}}}^n \binom{n}{k} a^k,$$

where $\binom{n}{k}$ is the binomial coefficient with the convention $\binom{n}{k} = 0$ for $k < 0$ or $k > n$.

Then we have the following theorem.

Theorem 2.2. Let $m, n \in \mathbb{Z}^+$, and $k \in \mathbb{Z}$. Write

$$W_n(k, m) = \sum_{\substack{l=1 \\ (l, m)=1}}^m \zeta_m^{-kl} (1 + a\zeta_m^l)^n,$$

and

$$A_m(x) = \prod_{\substack{l=1 \\ (l,m)=1}}^m (x - 1 - a\zeta_m^l) = \sum_{s=0}^{\varphi(m)} b_s x^s.$$

Then

$$A_m(x) \in \mathbb{Z}[x] \quad \text{and} \quad \sum_{s=0}^{\varphi(m)} b_s W_{n+s}(k, m) = 0.$$

Moreover, for any $r \in \mathbb{Z}$ we have

$$\left[\begin{matrix} n \\ r \end{matrix} \right]_m (a) = \frac{1}{m} \sum_{d|m} W_n(r, d).$$

Proof. It is easy to see that the coefficients of $A_m(x+1)$ are symmetric polynomials in those primitive m -th roots of unity with integer coefficients. Since

$$\Phi_m(x) = \prod_{\substack{l=1 \\ (l,m)=1}}^m (x - \zeta_m^l) \in \mathbb{Z}[x],$$

$A_m(x+1) \in \mathbb{Z}[x]$ by Fundamental Theorem on Symmetric Polynomials. Therefore $A_m(x) \in \mathbb{Z}[x]$.

For any positive integer n , we clearly have

$$\begin{aligned} \sum_{s=0}^{\varphi(m)} b_s W_{n+s}(k, m) &= \sum_{s=0}^{\varphi(m)} b_s \sum_{\substack{l=1 \\ (l,m)=1}}^m \zeta_m^{-kl} (1 + a\zeta_m^l)^{n+s} \\ &= \sum_{\substack{l=1 \\ (l,m)=1}}^m \zeta_m^{-kl} (1 + a\zeta_m^l)^n \sum_{s=0}^{\varphi(m)} b_s (1 + a\zeta_m^l)^s \\ &= \sum_{\substack{l=1 \\ (l,m)=1}}^m \zeta_m^{-kl} (1 + a\zeta_m^l)^n A_m(1 + a\zeta_m^l) \\ &= 0. \end{aligned}$$

Let $r \in \mathbb{Z}$, then we have

$$\begin{aligned}
\left[\begin{matrix} n \\ r \end{matrix} \right]_m (a) &= \sum_{k=0}^n \binom{n}{k} a^k \cdot \frac{1}{m} \sum_{l=1}^m \zeta_m^{(k-r)l} \\
&= \frac{1}{m} \sum_{l=1}^m \zeta_m^{-rl} (1 + a \zeta_m^l)^n \\
&= \frac{1}{m} \sum_{d|m} \sum_{\substack{b=1 \\ (b,d)=1}}^d \zeta_d^{-rb} (1 + a \zeta_d^b)^n \\
&= \frac{1}{m} \sum_{d|m} W_n(r, d).
\end{aligned}$$

This ends the proof. $\square$

Note that the theorem is a generalization of Theorem 1 of [13].

Remark 2.3. *The last result shows that $\left[\begin{matrix} n \\ r \end{matrix} \right]_m (a)$ can be expressed in terms of some linearly recurrent sequences with orders not exceeding $\varphi(m)$.*

Now we list $A_m(x)$ for $1 \leq m \leq 6$:

$$\begin{aligned}
A_1(x) &= x - 1 - a, A_2(x) = x - 1 + a, \\
A_3(x) &= x^2 - (2 - a)x + a^2 - a + 1, A_4(x) = x^2 - 2x + a^2 + 1, \\
A_5(x) &= x^4 - (4 - a)x^3 + (a^2 - 3a + 6)x^2 + (a^3 - 2a^2 + 3a + 4) + a^4 - a^3 + a^2 - a + 1, \\
A_6(x) &= x^2 - (a + 2)x + a^2 + a + 1.
\end{aligned}$$

Lemma 2.4. ([13]) *Let m, c be integers with $m > 0$. Then we have*

$$\sum_{d|m} \mu\left(\frac{m}{d}\right) d \delta_{d|c} = \varphi(m) \frac{\mu(m/(c, m))}{\varphi(m/(c, m))},$$

where

$$\delta_{d|c} = \begin{cases} 1, & \text{if } d \mid c \text{ holds;} \\ 0, & \text{otherwise.} \end{cases}$$

Proof. We can find that both sides are multiplicative with respect to m , thus we only need to prove it when m is a prime power. For any prime p and positive integer

k , we have

$$\begin{aligned}
\sum_{d|p^k} \mu\left(\frac{p^k}{d}\right) d \delta_{d|c} &= \sum_{s=0}^k \mu(p^{k-s}) p^s \delta_{p^s|c} \\
&= p^k \delta_{p^k|c} - p^{k-1} \delta_{p^{k-1}|c} \\
&= \begin{cases} p^k - p^{k-1} & \text{if } p^k \mid c, \\ -p^{k-1} & \text{if } p^{k-1} \parallel c, \\ 0 & \text{if } p^{k-1} \nmid c. \end{cases} \\
&= \varphi(p^k) \frac{\mu(p^k/(c, p^k))}{\varphi(p^k/(c, p^k))}.
\end{aligned}$$

This concludes the proof. $\square$

Theorem 2.5. *Let $m, n \in \mathbb{Z}^+, r \in \mathbb{Z}$. Then*

$$W_n(r, m) = \varphi(m) \sum_{k=0}^n \frac{\mu(m/(k-r, m))}{\varphi(m/(k-r, m))} \binom{n}{k} a^k.$$

Proof. By Theorem 2.2, Lemma 2.4 and Möbius Inversion Theorem, we have

$$\begin{aligned}
W_n(r, m) &= \sum_{d|m} \mu\left(\frac{m}{d}\right) d \left[\begin{matrix} n \\ r \end{matrix} \right]_d (a) \\
&= \sum_{d|m} \mu\left(\frac{m}{d}\right) d \sum_{k=0}^n \binom{n}{k} a^k \delta_{d|k-r} \\
&= \sum_{k=0}^n \binom{n}{k} a^k \sum_{d|m} \mu\left(\frac{m}{d}\right) d \delta_{d|k-r} \\
&= \varphi(m) \sum_{k=0}^n \frac{\mu(m/(k-r, m))}{\varphi(m/(k-r, m))} \binom{n}{k} a^k.
\end{aligned}$$

$\square$

Corollary 2.6. *Let m, n be two relatively prime positive integers. Then we have*

$$W_n(0, m) - \varphi(m) - \mu(m) a^n = \varphi(m) n \sum_{k=1}^{n-1} \frac{\mu(m/(m, k))}{\varphi(m/(m, k))} \binom{n-1}{k-1} \frac{a^k}{k}$$

and

$$W_n(n, m) - \varphi(m) a^n - \mu(m) = \varphi(m) n \sum_{k=1}^{n-1} \binom{n-1}{k-1} \frac{\mu(m/(m, k))}{\varphi(m/(m, k))} \frac{a^{n-k}}{k}.$$

Proof. Since $\binom{n}{k} = \frac{n}{k} \binom{n-1}{k-1}$ for $1 \leq k \leq n$, we can derive the results by setting $r = 0, n$ respectively in Theorem 2.5, $\square$

Corollary 2.7. *Let $m \in \mathbb{Z}^+$ and p be an odd prime not dividing am . Then we have*

$$\frac{W_p(0, m) - \varphi(m) - \mu(m)a^p}{p} \equiv -\varphi(m) \sum_{k=1}^{p-1} \frac{\mu(m/(m, k))}{\varphi(m/(m, k))} \cdot \frac{(-a)^k}{k} \pmod{p},$$

and

$$\frac{W_p(p, m) - \varphi(m)a^p - \mu(m)}{p} \equiv \varphi(m) \sum_{k=1}^{p-1} \frac{\mu(m/(m, k))}{\varphi(m/(m, k))} \cdot \frac{1}{k(-a)^{k-1}} \pmod{p}.$$

Proof. Since $\binom{p-1}{k} = (-1)^k$ for $0 \leq k \leq p-1$, the results follow from Corollary 2.6. $\square$

3. Some New Congruences

In this section, we give some new congruences by using the results of [16].

Lemma 3.1. *Let $p \nmid 3a(2-a)(a^3+1)$ be an odd prime, and $\{u_n\}_{n \geq 0}, \{v_n\}_{n \geq 0}$ be the Lucas sequences defined as*

$$u_0 = 0, u_1 = 1, u_{n+1} = (2-a)u_n - (a^2-a+1)u_{n-1} \text{ for } n \geq 1;$$

$$v_0 = 2, v_1 = (2-a), v_{n+1} = (2-a)v_n - (a^2-a+1)v_{n-1} \text{ for } n \geq 1.$$

Then we have:

(1)

$$\frac{u_p - \left(\frac{-3}{p}\right)}{p} \equiv \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^{k-1}}{2k-1} \cdot \left(\frac{a}{2-a}\right)^{2k-2} + \left(\frac{-3}{p}\right) \left(q_p(a) - q_p(2) + \frac{1}{2}q_p(3)\right) \pmod{p};$$

(2)

$$\frac{v_p - (2-a)}{p} \equiv (2-a) \left[-\frac{1}{2} \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^k}{k} \cdot \left(\frac{a}{2-a}\right)^{2k} - q_p(2) + q_p(2-a) \right] \pmod{p}.$$

Proof. By Lemmas 2.1 and 2.2 of [16], we have $u_p = \frac{1}{a\sqrt{-3}} \left[\left(\frac{2-a}{2} + \frac{a}{2}\sqrt{-3} \right)^p - \left(\frac{2-a}{2} - \frac{a}{2}\sqrt{-3} \right)^p \right]$, $v_p = \left(\frac{2-a}{2} + \frac{a}{2}\sqrt{-3} \right)^p + \left(\frac{2-a}{2} - \frac{a}{2}\sqrt{-3} \right)^p$, and $u_p \equiv \left(\frac{-3}{p} \right) \pmod{p}$, $v_p = (2-a)u_p - 2(a^2 - a + 1)u_{p-1} = 2u_{p+1} - (2-a)u_p \equiv (2-a) \pmod{p}$. Then

$$\begin{aligned} 2^{p-1}u_p &= \sum_{\substack{k=0 \\ k \text{ odd}}}^p \binom{p}{k} (2-a)^{p-k} (a\sqrt{-3})^{k-1} \\ &= a^{p-1}(-3)^{\frac{p-1}{2}} + \sum_{k=1}^{\frac{p-1}{2}} \binom{p}{2k-1} (2-a)^{p-2k+1} a^{2k-2} (-3)^{k-1} \\ &= a^{p-1}(-3)^{\frac{p-1}{2}} + p \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^{k-1}}{2k-1} \binom{p-1}{2k-2} (2-a)^{p-2k+1} a^{2k-2} \\ &\equiv a^{p-1}(-3)^{\frac{p-1}{2}} + p \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^{k-1}}{2k-1} \cdot \left(\frac{a}{2-a} \right)^{2k-2} \pmod{p^2}, \end{aligned}$$

and

$$\begin{aligned} 2^{p-1}v_p &= \sum_{\substack{k=0 \\ k \text{ even}}}^p \binom{p}{k} (2-a)^{p-k} (a\sqrt{-3})^k \\ &= (2-a)^p + \sum_{k=1}^{\frac{p-1}{2}} \binom{p}{2k} (2-a)^{p-2k} a^{2k} (-3)^k \\ &= (2-a)^p + p \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^k}{2k} \binom{p-1}{2k-1} (2-a)^{p-2k} a^{2k} \\ &\equiv (2-a)^p - \frac{2-a}{2} p \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^k}{k} \cdot \left(\frac{a}{2-a} \right)^{2k} \pmod{p^2}. \end{aligned}$$

Hence (1) and (2) follow from Lemma 2.6(1) of [16]. $\square$

Corollary 3.2. *Let $p \nmid 3a(2-a)(a^3+1)$ be an odd prime. Then we have*

$$\sum_{k=1}^{\left[\frac{p}{3}\right]} \frac{(-a)^{3k}}{k} \equiv (2-a) \left[\frac{1}{2} \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^k}{k} \cdot \left(\frac{a}{2-a} \right)^{2k} + q_p(2) - q_p(2-a) \right] - (a+1)q_p(a+1) \pmod{p}.$$

Proof. The result follows from Lemma 4.9 of [16] and Lemma 3.1(2). $\square$

Theorem 3.3. Let $p \nmid 3a(a-1)(2-a)(a^3+1)$ be an odd prime, and $\{u_n\}_{n \geq 0}$ be the Lucas sequence defined as

$$u_0 = 0, u_1 = 1, u_{n+1} = (2-a)u_n - (a^2-a+1)u_{n-1} \text{ for } n \geq 1.$$

(1) If $p \equiv 1 \pmod{3}$, we have

$$\frac{u_{p-1}}{p} \equiv -\frac{2}{a(a-1)} \sum_{k=1}^{\frac{p-1}{3}} \frac{(-a)^{3k-1}}{3k-1} + \frac{a+1}{3a(a-1)} (q_p(a^2-a+1) - 2q_p(a+1)) \pmod{p}$$

and

$$\begin{aligned} \sum_{k=1}^{\frac{p-1}{3}} \frac{(-a)^{3k-1}}{3k-1} &\equiv \frac{a(a-1)}{a-2} \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^{k-1}}{2k-1} \cdot \left(\frac{a}{2-a}\right)^{2k-2} \\ &\quad + \frac{a(a-1)}{a-2} [q_p(a) - q_p(2) + \frac{1}{2}q_p(3)] \\ &\quad - \frac{1}{3}(a+1)q_p(a+1) - \frac{a^2-a+1}{3(a-2)}q_p(a^2-a+1) \pmod{p}. \end{aligned}$$

(2) If $p \equiv 2 \pmod{3}$, we have

$$\frac{u_{p+1}}{p} \equiv \frac{2(a^2-a+1)}{a(a-1)} \sum_{k=1}^{\frac{p+1}{3}} \frac{(-a)^{3k-2}}{3k-2} - \frac{a^3+1}{3a(a-1)} (q_p(a^2-a+1) - 2q_p(a+1)) \pmod{p}$$

and

$$\begin{aligned} \sum_{k=1}^{\frac{p+1}{3}} \frac{(-a)^{3k-2}}{3k-2} &\equiv -\frac{a(a-1)}{a-2} \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^{k-1}}{2k-1} \cdot \left(\frac{a}{2-a}\right)^{2k-2} \\ &\quad + \frac{a(a-1)}{a-2} [q_p(a) - q_p(2) + \frac{1}{2}q_p(3)] \\ &\quad - \frac{1}{3}(a+1)q_p(a+1) - \frac{a^2-a+1}{3(a-2)}q_p(a^2-a+1) \pmod{p}. \end{aligned}$$

Proof. Since $(2-a)^2 - 4(a^2-a+1) = -3a^2$, we have $p \mid u_{p-\left(\frac{-3}{p}\right)}$ by Lemma 2.2 of [16]. Let $\{v_n\}_{n \geq 0}$ be the Lucas sequence define as Lemma 3.1.

(1) By Lemma 2.1 and Theorem 4.1 of [16], we have $-(a+1)u_p + (a^2-a+1)u_{p-1} = 3 \left[\begin{smallmatrix} p \\ 2 \end{smallmatrix} \right]_3 (a) - (1+a)^p$ and $v_{p-1} = 2u_p - (2-a)u_{p-1}$. Thus by Lemma 2.4 of [16], we have

$$\begin{aligned} 3a(a-1)u_{p-1} &= 6 \left[\begin{smallmatrix} p \\ 2 \end{smallmatrix} \right]_3 (a) - 2(1+a)^p + (a+1)v_{p-1} \\ &\equiv -6p \sum_{k=1}^{\frac{p-1}{3}} \frac{(-a)^{3k-1}}{3k-1} + (a+1) [(v_p - 2) - 2((a+1)^{p-1} - 1)] \pmod{p^2} \end{aligned}$$

and

$$\begin{aligned}
3a(a-1)(u_p-1) &= 3(2-a) \left[\begin{matrix} p \\ 2 \end{matrix} \right]_3 (a) - (2-a)(1+a)^p + (a^2-a+1)v_{p-1} - 3a(a-1) \\
&\equiv -3(2-a)p \sum_{k=1}^{\frac{p-1}{3}} \frac{(-a)^{3k-1}}{3k-1} - (2-a)(1+a)((a+1)^{p-1}-1) \\
&\quad + (a^2-a+1)(v_{p-1}-2) \pmod{p^2}.
\end{aligned}$$

Thence by Lemma 2.7 of [16] and Lemma 3.1(1),

$$\frac{u_{p-1}}{p} \equiv -\frac{2}{a(a-1)} \sum_{k=1}^{\frac{p-1}{3}} \frac{(-a)^{3k-1}}{3k-1} + \frac{a+1}{3a(a-1)} (q_p(a^2-a+1) - 2q_p(a+1)) \pmod{p},$$

and

$$\begin{aligned}
\sum_{k=1}^{\frac{p-1}{3}} \frac{(-a)^{3k-1}}{3k-1} &\equiv \frac{a(a-1)}{a-2} \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^{k-1}}{2k-1} \cdot \left(\frac{a}{2-a} \right)^{2k-2} \\
&\quad + \frac{a(a-1)}{a-2} [q_p(a) - q_p(2) + \frac{1}{2}q_p(3)] \\
&\quad - \frac{1}{3}(a+1)q_p(a+1) - \frac{a^2-a+1}{3(a-2)}q_p(a^2-a+1) \pmod{p}.
\end{aligned}$$

(2) By Lemma 2.1 and Theorem 4.1 of [16], we have $-u_{p+1} + (a+1)u_p = 3 \left[\begin{matrix} p \\ 1 \end{matrix} \right]_3 (a) - (1+a)^p$ and $v_{p+1} = (2-a)u_{p+1} - 2(a^2-a+1)u_p$. Thus by Lemmas 2.4 of [16], we have

$$\begin{aligned}
-3a(a-1)u_{p+1} &= 6(a^2-a+1) \left[\begin{matrix} p \\ 1 \end{matrix} \right]_3 (a) - 2(a^2-a+1)(1+a)^p + (a+1)v_{p+1} \\
&\equiv -6(a^2-a+1)p \sum_{k=1}^{\frac{p+1}{3}} \frac{(-a)^{3k-2}}{3k-2} - 2(a^2-a+1)(a+1)[(a+1)^{p-1}-1] \\
&\quad + (a+1)[v_{p+1} - 2(a^2-a+1)] \pmod{p^2}
\end{aligned}$$

and

$$\begin{aligned}
-3a(a-1)(u_p+1) &= 3(2-a) \left[\begin{matrix} p \\ 1 \end{matrix} \right]_3 (a) - (2-a)(1+a)^p + v_{p+1} - 3a(a-1) \\
&\equiv -3(2-a)p \sum_{k=1}^{\frac{p+1}{3}} \frac{(-a)^{3k-2}}{3k-2} - (2-a)(a+1)[(a+1)^{p-1}-1] \\
&\quad + v_{p+1} - 2(a^2-a+1) \pmod{p^2}.
\end{aligned}$$

Thence by Lemma 2.7 of [16] and Lemma 3.1(1),

$$\frac{u_{p+1}}{p} \equiv \frac{2(a^2 - a + 1)}{a(a-1)} \sum_{k=1}^{\frac{p+1}{3}} \frac{(-a)^{3k-2}}{3k-2} - \frac{a^3 + 1}{3a(a-1)} (q_p(a^2 - a + 1) - 2q_p(a+1)) \pmod{p},$$

and

$$\begin{aligned} \sum_{k=1}^{\frac{p+1}{3}} \frac{(-a)^{3k-2}}{3k-2} &\equiv -\frac{a(a-1)}{a-2} \sum_{k=1}^{\frac{p-1}{2}} \frac{(-3)^{k-1}}{2k-1} \cdot \left(\frac{a}{2-a}\right)^{2k-2} \\ &\quad + \frac{a(a-1)}{a-2} [q_p(a) - q_p(2) + \frac{1}{2}q_p(3)] \\ &\quad - \frac{1}{3}(a+1)q_p(a+1) - \frac{a^2 - a + 1}{3(a-2)}q_p(a^2 - a + 1) \pmod{p}. \end{aligned}$$

□

Set $a = -2$ in Corollary 3.2 and Theorem 3.3, we have the following two corollaries.

Corollary 3.4. *Let $p \neq 3, 7$ be and odd prime. Then we have*

(1)

$$\sum_{k=1}^{\lfloor \frac{p}{3} \rfloor} \frac{8^k}{k} \equiv \sum_{k=1}^{\frac{p-1}{2}} \frac{2}{k} \cdot \left(-\frac{3}{4}\right)^k - 4q_p(2) \pmod{p}.$$

(2) *If $p \equiv 1 \pmod{3}$,*

$$\sum_{k=1}^{\frac{p-1}{3}} \frac{8^k}{3k-1} \equiv 4 \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{2k-1} \cdot \left(-\frac{3}{4}\right)^k - \frac{3}{2}q_p(3) + \frac{7}{6}q_p(7) \pmod{p}.$$

If $p \equiv 2 \pmod{3}$,

$$\sum_{k=1}^{\frac{p+1}{3}} \frac{8^k}{3k-2} \equiv -8 \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{2k-1} \cdot \left(-\frac{3}{4}\right)^k - 3q_p(2) + \frac{7}{3}q_p(7) \pmod{p}.$$

Corollary 3.5. *Let $p \neq 3, 7$ be and odd prime, and $\{u_n\}_{n \geq 0}$ be the Lucas sequence defined as*

$$u_0 = 0, u_1 = 1, u_{n+1} = 4u_n - 7u_{n-1} \text{ for } n \geq 1.$$

Then, if $p \equiv 1 \pmod{3}$,

$$\frac{u_{p-1}}{p} \equiv -\frac{1}{6} \sum_{k=1}^{\frac{p-1}{3}} \frac{8^k}{3k-1} - \frac{1}{18}q_p(7) \pmod{p},$$

if $p \equiv 2 \pmod{3}$,

$$\frac{u_{p+1}}{p} \equiv \frac{7}{12} \sum_{k=1}^{\frac{p+1}{3}} \frac{8^k}{3k-2} + \frac{7}{18} q_p(7) \pmod{p}.$$

The following theorem can reduce the summation terms occurring in the expression of Lucas quotients in Corollary 4.11 of [16] and Corollary 3.5.

Theorem 3.6. *Let $p \neq 3, 7$ be an odd prime, and $\{u_n\}_{n \geq 0}$ be the Lucas sequence defined as Corollary 3.5. Then if $p \equiv 1 \pmod{3}$,*

$$\begin{aligned} \frac{u_{p-1}}{p} &\equiv \frac{1}{6} \sum_{k=1}^{\frac{p-1}{6}} \frac{64^k}{k} + \frac{1}{3} q_p(7) + \frac{1}{2} q_p(3) \\ &\equiv -\frac{1}{3} \sum_{k=1}^{\frac{p-1}{6}} \frac{64^k}{6k-1} - \frac{1}{18} q_p(7) + \frac{1}{6} q_p(3) \pmod{p}, \end{aligned}$$

if $p \equiv 2 \pmod{3}$,

$$\begin{aligned} \frac{u_{p+1}}{p} &\equiv -\frac{7}{6} \sum_{k=1}^{\frac{p-5}{6}} \frac{64^k}{k} - \frac{7}{3} q_p(7) - \frac{7}{2} q_p(3) \\ &\equiv \frac{7}{6} \sum_{k=1}^{\frac{p+1}{6}} \frac{64^k}{6k-2} + \frac{7}{18} q_p(7) + \frac{7}{6} q_p(3) \pmod{p}. \end{aligned}$$

Proof. By Lemma 2.4 and Theorem 4.5 of [16], if $p \equiv 1 \pmod{3}$,

$$3^{p-1} - (-3)^{\frac{p-1}{2}} = \left[\begin{matrix} p \\ 2 \end{matrix} \right]_3 (2) \equiv -p \sum_{k=1}^{\frac{p-1}{3}} \frac{(-2)^{3k-1}}{3k-1} \pmod{p^2},$$

if $p \equiv 2 \pmod{3}$,

$$3^{p-1} + (-3)^{\frac{p-1}{2}} = \left[\begin{matrix} p \\ 1 \end{matrix} \right]_3 (2) \equiv -p \sum_{k=1}^{\frac{p+1}{3}} \frac{(-2)^{3k-2}}{3k-2} \pmod{p^2}.$$

Thus by the Lemma 2.6 of [16], we have

$$\sum_{k=1}^{\frac{p-1}{3}} \frac{(-8)^k}{3k-1} \equiv q_p(3) \pmod{p} \text{ if } p \equiv 1 \pmod{3},$$

and

$$\sum_{k=1}^{\frac{p+1}{3}} \frac{(-8)^k}{3k-2} \equiv -2q_p(3) \pmod{p} \text{ if } p \equiv 2 \pmod{3}.$$

Hence the results follow from Corollaries 4.7 and 4.11 of [16] and Corollary 3.5. $\square$

4. A Specific Lucas Sequence

Let $A, B \in \mathbb{Z}$. The Lucas sequences $u_n = u_n(A, B) (n \in \mathbb{N})$ and $v_n = v_n(A, B) (n \in \mathbb{N})$ are defined by

$$u_0 = 1, u_1 = 1, u_{n+1} = Bu_n - Au_{n-1} (n \geq 1);$$

$$v_0 = 2, v_1 = B, v_{n+1} = Bv_n - Av_{n-1} (n \geq 1).$$

Next we give some properties of the Lucas sequences with $A = 5$ and $B = 2$. We need some lemmas. Let $D = B^2 - 4A$.

Lemma 4.1. *Let p be an odd prime not dividing DA .*

- (1) *If $p \equiv 1 \pmod{4}$, then $p \mid u_{\frac{p-1}{4}}$ if and only if $v_{\frac{p-1}{2}} \equiv 2A^{\frac{p-1}{4}} \pmod{p}$ and $p \mid v_{\frac{p-1}{4}}$ if and only if $v_{\frac{p-1}{2}} \equiv -2A^{\frac{p-1}{4}} \pmod{p}$.*
- (2) *If $p \equiv 3 \pmod{4}$, then $p \mid u_{\frac{p+1}{4}}$ if and only if $v_{\frac{p+1}{2}} \equiv 2A^{\frac{p+1}{4}} \pmod{p}$ and $p \mid v_{\frac{p+1}{4}}$ if and only if $v_{\frac{p+1}{2}} \equiv -2A^{\frac{p+1}{4}} \pmod{p}$.*

Proof. (1) and (2) follow from the fact that $v_{2n} = v_n^2 - 2A^n = Du_n^2 + 2A^n$. $\square$

Lemma 4.2. ([9]) *Let p be an odd prime and A' be an integer such that $4A' \equiv B^2 - 4A \pmod{p}$. Let $u'_n = u_n(A', B)$, $v'_n = v_n(A', B)$. Then we have*

$$u_{\frac{p+1}{2}} \equiv \frac{1}{2} \left(\frac{2}{p} \right) v'_{\frac{p-1}{2}} \pmod{p}, \quad u_{\frac{p-1}{2}} \equiv - \left(\frac{2}{p} \right) u'_{\frac{p-1}{2}} \pmod{p},$$

$$v_{\frac{p+1}{2}} \equiv \left(\frac{2}{p} \right) v'_{\frac{p+1}{2}} \pmod{p}, \quad v_{\frac{p-1}{2}} \equiv 2 \left(\frac{2}{p} \right) u'_{\frac{p+1}{2}} \pmod{p}.$$

Remark 4.3. (1) *Let $S_n = u_n(1, 4)$, $T_n = v_n(1, 4)$. For any prime $p > 3$, by the facts that $u'_n = u_n(3, 4) = \frac{1}{2}(3^n - 1)$ and $v'_n = v_n(3, 4) = 3^n + 1$, we have*

$$S_{\frac{p+1}{2}} \equiv \frac{1}{2} \left(\frac{2}{p} \right) \left[\left(\frac{3}{p} \right) + 1 \right] \pmod{p}, \quad S_{\frac{p-1}{2}} \equiv -\frac{1}{2} \left(\frac{2}{p} \right) \left[\left(\frac{3}{p} \right) - 1 \right] \pmod{p},$$

$$T_{\frac{p+1}{2}} \equiv \left(\frac{2}{p} \right) \left[3 \left(\frac{3}{p} \right) + 1 \right] \pmod{p}, \quad T_{\frac{p-1}{2}} \equiv \left(\frac{2}{p} \right) \left[3 \left(\frac{3}{p} \right) - 1 \right] \pmod{p}.$$

Thus by Lemma 4.1, $p \mid S_{\frac{p+1}{4}}$ iff $p \equiv 1, 19 \pmod{24}$ and $p \mid T_{\frac{p+1}{4}}$ iff $p \equiv 7, 13 \pmod{24}$. Sun [13] got these by studying the sum (1) for $a = 1$ and $m = 12$;

- (2) Let $P_n = u_n(-1, 2)$, $Q_n = v_n(-1, 2)$ and $u'_n = u_n(2, 2)$, $v_n = v'_n(2, 2)$. For any odd prime p , by the facts that $u'_{4n} = 0, u'_{4n+1} = (-4)^n, u'_{4n+2} = u'_{4n+3} = 2(-4)^n$ and $v'_{4n} = v'_{4n+1} = 2(-4)^n, u'_{4n+2} = 0, u'_{4n+3} = (-4)^{n+1}$, we have

$$P_{\frac{p-(\frac{2}{p})}{2}} \equiv \begin{cases} 0 \pmod{p}, & \text{if } p \equiv 1 \pmod{4}, \\ (-1)^{[\frac{p+5}{8}]} 2^{\frac{p-3}{4}} \pmod{p}, & \text{if } p \equiv 3 \pmod{4}, \end{cases}$$

$$Q_{\frac{p-(\frac{2}{p})}{2}} \equiv \begin{cases} (-1)^{[\frac{p}{8}]} 2^{\frac{p+3}{4}} \pmod{p}, & \text{if } p \equiv 1 \pmod{4}, \\ 0 \pmod{p}, & \text{if } p \equiv 3 \pmod{4}, \end{cases}$$

and

$$P_{\frac{p+(\frac{2}{p})}{2}} \equiv (-1)^{[\frac{p+1}{8}]} 2^{[\frac{p}{4}]} \pmod{p}, \quad Q_{\frac{p+(\frac{2}{p})}{2}} \equiv (-1)^{[\frac{p+5}{8}]} 2^{[\frac{p+5}{4}]} \pmod{p}.$$

Sun got [7] these by studying the sum (1) for $a = 1$ and $m = 8$.

Lemma 4.4. Let $p \nmid B$ be an odd prime and A' be an integer such that $A' \equiv \frac{A}{B^2} \pmod{p}$. Let $u'_n = u_n(A', 1)$, $v'_n = v_n(A', 1)$. Then we have

$$\begin{aligned} u_{\frac{p+1}{2}} &\equiv \left(\frac{B}{p}\right) u'_{\frac{p+1}{2}} \pmod{p}, & u_{\frac{p-1}{2}} &\equiv \frac{1}{B} \left(\frac{B}{p}\right) u'_{\frac{p-1}{2}} \pmod{p}, \\ v_{\frac{p+1}{2}} &\equiv B \left(\frac{B}{p}\right) v'_{\frac{p+1}{2}} \pmod{p}, & v_{\frac{p-1}{2}} &\equiv \left(\frac{B}{p}\right) v'_{\frac{p-1}{2}} \pmod{p}. \end{aligned}$$

Proof. By Lemma 2.1 of [16] and $D' = 1 - 4A' \equiv \frac{D}{B^2} \pmod{p}$, we have

$$\begin{aligned} u_n &= 2 \sum_{\substack{k=0 \\ k \text{ odd}}}^n \binom{n}{k} \left(\frac{B}{2}\right)^{n-k} \left(\frac{D}{2}\right)^{\frac{k-1}{2}} \\ &= 2B^{n-1} \sum_{\substack{k=0 \\ k \text{ odd}}}^n \binom{n}{k} \left(\frac{1}{2}\right)^{\frac{p-1}{2}-k} \left(\frac{D}{2B^2}\right)^{\frac{k-1}{2}} \\ &\equiv 2B^{n-1} \sum_{\substack{k=0 \\ k \text{ odd}}}^n \binom{n}{k} \left(\frac{1}{2}\right)^{n-k} \left(\frac{D'}{2}\right)^{\frac{k-1}{2}} \\ &= B^{n-1} u'_n \pmod{p}, \end{aligned}$$

and

$$\begin{aligned}
v_n &= 2 \sum_{\substack{k=0 \\ k \text{ even}}}^n \binom{n}{k} \left(\frac{B}{2}\right)^{n-k} \left(\frac{D}{2}\right)^{\frac{k}{2}} \\
&= 2B^n \sum_{\substack{k=0 \\ k \text{ even}}}^n \binom{n}{k} \left(\frac{1}{2}\right)^{\frac{p-1}{2}-k} \left(\frac{D}{2B^2}\right)^{\frac{k}{2}} \\
&\equiv 2B^n \sum_{\substack{k=0 \\ k \text{ even}}}^n \binom{n}{k} \left(\frac{1}{2}\right)^{n-k} \left(\frac{D'}{2}\right)^{\frac{k}{2}} \\
&= B^n v'_n \pmod{p},
\end{aligned}$$

Thus

$$\begin{aligned}
u_{\frac{p+1}{2}} &\equiv B^{\frac{p-1}{2}} u'_{\frac{p+1}{2}} \equiv \left(\frac{B}{p}\right) u'_{\frac{p+1}{2}} \pmod{p}, \\
u_{\frac{p-1}{2}} &\equiv B^{\frac{p-3}{2}} u'_{\frac{p-1}{2}} \equiv \frac{1}{B} \left(\frac{B}{p}\right) u'_{\frac{p-1}{2}} \pmod{p}, \\
v_{\frac{p+1}{2}} &\equiv B^{\frac{p+1}{2}} v'_{\frac{p+1}{2}} \equiv B \left(\frac{B}{p}\right) v'_{\frac{p+1}{2}} \pmod{p}, \\
v_{\frac{p-1}{2}} &\equiv B^{\frac{p-1}{2}} v'_{\frac{p-1}{2}} \equiv \left(\frac{B}{p}\right) v'_{\frac{p-1}{2}} \pmod{p}.
\end{aligned}$$

□

Theorem 4.5. *Let $p \neq 5$ be an odd prime and $\{U_n\}_{n \geq 0}$ and $\{V_n\}_{n \geq 0}$ be the Lucas sequences defined as*

$$U_0 = 0, U_1 = 1, U_{n+1} = 2U_n - 5U_{n-1} \text{ for } n \geq 1;$$

$$V_0 = 2, V_1 = 2, V_{n+1} = 2V_n - 5V_{n-1} \text{ for } n \geq 1.$$

(1) *If $p \equiv \pm 1 \pmod{5}$, we have*

$$\begin{aligned}
U_{\frac{p+(-1)}{2}} &\equiv \left(\frac{-1}{p}\right) (-1)^{[\frac{p+5}{10}]} 5^{[\frac{p}{4}]} \pmod{p}, \\
U_{\frac{p-(-1)}{2}} &\equiv 0 \pmod{p}, \\
V_{\frac{p+(-1)}{2}} &\equiv 2(-1)^{[\frac{p+5}{10}]} 5^{[\frac{p}{4}]} \pmod{p}, \\
V_{\frac{p-(-1)}{2}} &\equiv 2(-1)^{[\frac{p+5}{10}]} 5^{[\frac{p+1}{4}]} \pmod{p}.
\end{aligned}$$

(2) If $p \equiv \pm 2 \pmod{5}$, we have

$$\begin{aligned} U_{\frac{p+(\frac{-1}{p})}{2}} &\equiv \frac{1}{2} \left(\frac{-1}{p} \right) (-1)^{[\frac{p+5}{10}]} 5^{[\frac{p}{4}]} \pmod{p}, \\ U_{\frac{p-(\frac{-1}{p})}{2}} &\equiv \frac{1}{2} \left(\frac{-1}{p} \right) (-1)^{[\frac{p+5}{10}]} 5^{[\frac{p+1}{4}]} \pmod{p}, \\ V_{\frac{p+(\frac{-1}{p})}{2}} &\equiv 4(-1)^{[\frac{p-5}{10}]} 5^{[\frac{p}{4}]} \pmod{p}, \\ V_{\frac{p-(\frac{-1}{p})}{2}} &\equiv 0 \pmod{p}. \end{aligned}$$

Proof. Let $F_n = u_n(-1, 1)$ and $L_n = v_n(-1, 1)$ be Fibonacci sequence and its companion. Then by Lemmas 4.2 and 4.4, we have

$$\begin{aligned} U_{\frac{p+1}{2}} &\equiv \frac{1}{2} L_{\frac{p-1}{2}} \pmod{p}, & U_{\frac{p-1}{2}} &\equiv -\frac{1}{2} F_{\frac{p-1}{2}} \pmod{p}, \\ V_{\frac{p+1}{2}} &\equiv 2L_{\frac{p+1}{2}} \pmod{p}, & V_{\frac{p-1}{2}} &\equiv 2F_{\frac{p+1}{2}} \pmod{p}. \end{aligned}$$

Thus by Corollaries 1 and 2 of [11], we can derive the results. $\square$

Remark 4.6. In [16], we gave some congruences for the Lucas quotient $U_{p-(\frac{-1}{p})}/p$ by studying the sum (1) for $a = -2$ and $m = 4$.

Corollary 4.7. Let $p \neq 5$ be an odd prime, $\{U_n\}_{n \geq 0}$ and $\{V_n\}_{n \geq 0}$ be Lucas sequences defined as above.

- (1) If $p \equiv 1 \pmod{4}$, then $p \mid U_{\frac{p-1}{4}}$ if and only if $p \equiv 1 \pmod{20}$ and $p \mid V_{\frac{p-1}{4}}$ if and only if $p \equiv 9 \pmod{20}$.
- (2) If $p \equiv 3 \pmod{4}$, then $p \mid U_{\frac{p+1}{4}}$ if and only if $p \equiv 19 \pmod{20}$ and $p \mid V_{\frac{p+1}{4}}$ if and only if $p \equiv 11 \pmod{20}$.

Proof. (1) and (2) follow from Lemma 4.1 and Theorem 4.5. $\square$

Acknowledgments The work of this paper was supported by the NNSF of China (Grant No. 11471314), and the National Center for Mathematics and Interdisciplinary Sciences, CAS.

References

- [1] P.Bundschuh, C.Ji and Z.Shao, *A remarkable class of congruences*, Acta Sci. Math. (Szeged) **67** (2001), no. 3-4, 493-500.

- [2] R. Crandall and C. Pomerance, *Prime Numbers, A Computational Perspective*, Second edition, Springer, New York, 2005.
- [3] Y. Deng and Y. Pan, *The Sum of Binomial Coefficients and Integer Factorization*, *Integers* **16**(2016), Paper No. A42, 18pp.
- [4] R. L. Rivest, A. Shamir and L. Adleman, *A Method for Obtaining Digital Signatures and Public-key Cryptosystems*, *Comm. ACM* **21**(1978), 120–126.
- [5] Z. Shan and E.T.H. Wang, *A simple proof of a curious congruence by Sun*, *Proc. Amer. Math. Soc.* **127** (1999), no. 5, 1289–1291.
- [6] Z.-H. Sun, *Combinatorial Sum $\sum_{k=0, k \equiv r \pmod{m}}^n \binom{n}{k}$ and its Applications in Number Theory (I)*, *Nanjing Daxue Xuebao Shuxue Bannian Kan* **9** (1992), 227–240.
- [7] Z.-H. Sun, *Combinatorial Sum $\sum_{k=0, k \equiv r \pmod{m}}^n \binom{n}{k}$ and its Applications in Number Theory (II)*, *Nanjing Daxue Xuebao Shuxue Bannian Kan* **10**(1993), 105–118.
- [8] Z.-H. Sun, *Combinatorial Sum $\sum_{k=0, k \equiv r \pmod{m}}^n \binom{n}{k}$ and its Applications in Number Theory (III)*, *Nanjing Daxue Xuebao Shuxue Bannian Kan* **12**(1995), 90–102.
- [9] Z.H. Sun, *Values of Lucas Sequences Modulo Primes*, *The Rocky Mountain Journal of Mathematics.* **33**(2003), No.3,1123–1145.
- [10] Z.H. Sun, *Five Congruences for Primes*, *The Fibonacci Quartely.* **40**(2002), No.4,345–351.
- [11] Z.-H. Sun and Z.-W. Sun, *Fibonacci Numbers and Fermat's Last Theorem*, *Acta Arith.* **60**(1992), 371–388.
- [12] Z.-W. Sun, *A Congruence for Primes*, *Proc. Amer. Math. Soc.* **123**(1995), 1341–1346.
- [13] Z.-W. Sun, *On the Sum $\sum_{k \equiv r \pmod{m}} \binom{n}{k}$ and Related Congruences*, *Israel J. Math.* **128**(2002), 135–156.
- [14] D. D. Wall, *Fibonacci Series Modulo m* , *Amer. Math. Monthly* **67**(1960), 525–532.
- [15] H. C. Williams, *A Note on the Fibonacci Quotient $F_{p-\varepsilon}/p$* , *Canad. Math. Bull.* **25**(1982), 366–370.
- [16] J. Yang and Y. Deng, *Combinatorial Sums $\sum_{k \equiv r \pmod{m}} \binom{n}{k} a^k$ and Lucas Quotients*, arXiv:1606.08245 [math.NT].